



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

- First,  $3x+2z=3[3y+2u]... (1);$   
 Second,  $2x+6v=8\{2[1-(x+y)]+6[1-(v+w)]\}... (2);$   
 Third,  $2u+3w=5\{2[1-(z+u)]+3[1-(v+w)]\}... (3);$   
 Fourth,  $x+2z+3v=2(y+2u+3w)... (4);$   
 Fifth,  $x+z : y+u :: 11 : 5... (5);$  and  
 Sixth,  $6x+5z+2v : 13 :: 17 : 24... (6).$

By elimination, we have  $x=1$ ,  $y=0$ ,  $1-(x+y)=0$ ,  $z=\frac{3}{8}$ ,  $u=\frac{5}{8}$ ,  $1-(z+u)=0$ ,  $v=\frac{9}{8}$ ,  $w=\frac{5}{4}$ , and  $1-(v+w)=\frac{1}{8}$ .

We interpret these results as follows: That the first ingot was pure gold; that the second ingot was 9-carat gold, and 15-carat silver; and that the third ingot was 16-carat gold, 5-carat silver, and 3-carat copper.

Also solved in the same manner by G. B. M. Zerr, S. A. Corey, L. E. Newcomb, and G. W. Greenwood.

274. Proposed by R. D. CARMICHAEL, Anniston, Ala.

Find the limit of  $\frac{3^2+1}{3^2-1} \frac{5^2+1}{5^2-1} \frac{7^2+1}{7^2-1} \frac{11^2+1}{11^2-1} \dots$  where the squared numbers are the natural odd *primes* in order.

Solution by G. B. M. ZERR, Ph. D., Parsons, W. Va., and J. SCHEFFER, A. M., Kee Mar College, Hagerstown, Md.

Putting the expression in the form

$$\frac{(1+1/3^2)(1+1/5^2)(1+1/7^2)\dots}{(1-1/3^2)(1-1/5^2)(1-1/7^2)\dots} = \frac{s}{s'}$$

and remembering that  $(1+1/2^2)s = \frac{15}{\pi^2}$  and  $(1-1/2^2)s' = \frac{6}{\pi^2}$ , (pp. 133-134, Vol. V, No. 5), we have  $s/s' = 1\frac{1}{2}$ .

C. N. Schmall gives the following arithmetical solution of 269. When the boats first meet, combined distance traveled is equal to width of river; when they meet for the second time the distance traveled is equal to three times the width of river and that each boat has gone three times as far as when they first met. Hence one has gone  $3 \times 720$  yards = 2160 yards and has made one trip and 400 yards of the return trip. Hence, width of river = 2160 yards - 400 yards = 1760 yards = 1 mile.

## GEOMETRY.

302. Proposed by F. H. SAFFORD, Ph. D., University of Pennsylvania.

Through a given point within a circle draw any two chords, also a radius and a secant perpendicular to the radius. Let the extremities of the chords be taken as the vertices of a quadrilateral. Show that the sides of the quadrilateral, produced when necessary, cut the secant in points equidistant, in pairs, from the given point. [A proof by Euclidean geometry is preferred, as the problem was originally given to a high school class.] Must the given points be within the circle?